

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	NA

Hadlima (adalimumab-bwwd)

Override(s)	Approval Duration
Prior Authorization Quantity Limit	1 year

Medications	Quantity Limit
Hadlima (adalimumab-bwwd) 40 mg/0.4 mL PushTouch Autoinjector ^{##¥^S†‡}	2 syringes per 28 days
Hadlima (adalimumab-bwwd) 40 mg/0.8 mL prefilled syringe ^{##¥^S†‡}	2 syringes per 28 days

Override Criteria

[#]In the treatment of Rheumatoid Arthritis (RA): May approve up to four (4) syringes (40mg/0.8ml) (up to an additional two (2) syringes) every 28 days if the individual is unable to take concomitant methotrexate.

^{*}Initiation of therapy for adult Crohn's Disease (CD) or Ulcerative Colitis (UC): May approve up to four (4) additional syringes (40 mg/0.8 mL) in the first month (28 days) of treatment.

[¥]In the treatment of CD or UC: May approve up to an additional 2 (two) syringes (40 mg/0.8 mL) every 28 days if the individual has an inadequate response to standard maintenance dosing.

[^]Initiation of therapy for Plaque Psoriasis (Ps): May approve one (1) Ps starter pack **OR** up to two (2) additional syringes (40 mg/0.8 mL) in the first month (28 days) of treatment.

[§]Initiation of therapy for adult Hidradenitis Suppurativa (HS): May approve 1 (one) Crohn's Disease/Ulcerative Colitis/Hidradenitis Suppurativa starter pack **OR** up to 4 (four) additional pens or syringes (40 mg) in the first month (28 days) of treatment. Maintenance therapy: May approve up to 2 (two) additional pens or syringes (40 mg) per each 28 days.

[†]Initiation of therapy for pediatric Crohn's Disease (CD): Depending on individual's weight, may approve one (1) pediatric or adult Crohn's Disease starter pack **OR** up to 4 (four) additional pens or syringes (40 mg) in the first month (28 days) of treatment.

[‡]Initiation of therapy for Uveitis (UV): May approve up to 2 (two) additional pens or syringes (40 mg) in the first month (28 days) of treatment.

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	NA

APPROVAL CRITERIA

Requests for Hadlima (adalimumab-bwwd) may be approved for the following:

- I. Crohn's disease (CD) when each of the following criteria are met:
- A. Individual is 6 years of age or older with moderate to severe CD; **AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as 5-Aminosalicylic acid products, systemic corticosteroids, or immunosuppressants);
- AND**
- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to ONE (1) preferred biologic agent [Current preferred biologics include – Humira (adalimumab), Inflectra (infliximab-dyyb), Renflexis (infliximab-abda)] unless the following criteria are met:
 - 1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

- II. Ulcerative colitis (UC) when each of the following criteria are met:
- A. Individual is 18 years of age or older with moderate to severe UC; **AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as 5-Aminosalicylic acid products, systemic corticosteroids, or immunosuppressants);
- AND**
- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to ONE (1) preferred biologic agent [Current preferred biologics include – Humira (adalimumab), Inflectra (infliximab-dyyb), Renflexis (infliximab-abda)] unless the following criteria are met:
 - 1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

- III. Rheumatoid arthritis (RA) when each of the following criteria are met:
- A. Individual must be 18 years of age or older with moderate to severe RA; **AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [nonbiologic disease modifying anti-rheumatic agents (DMARDs) (such as methotrexate, sulfasalazine, leflunomide, or hydroxychloroquine)] (ACR 2015);
- AND**

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	NA

- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred agents [Current preferred agents include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:
1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

- IV. Ankylosing spondylitis (AS) when each of the following criteria are met:
- A. Individual is 18 years of age or older with moderate to severe AS; **AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [such as NSAIDs or nonbiologic disease modifying anti-rheumatic drugs (DMARDs) (such as sulfasalazine)];

AND

- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:
1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

- V. Polyarticular juvenile idiopathic arthritis (PJIA) when each of the following criteria are met:
- A. Individual is 2 years of age or older with moderate to severe PJIA; **AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [nonbiologic disease modifying anti-rheumatic drugs (DMARDs) (such as methotrexate)] (ACR 2019);

AND

- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:
1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

- VI. Psoriatic arthritis (PsA) when each of the following criteria are met:
- A. Individual is 18 years of age or older with moderate to severe PsA; **AND**
 - B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [nonbiologic disease modifying anti-

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	NA

rheumatic drugs (DMARDS) (such as methotrexate, sulfasalazine, or leflunomide)];

AND

C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:

1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

VII. Plaque psoriasis (Ps) (Psoriasis vulgaris) when each of the following criteria are met:

A. Individual is 18 years of age or older with

chronic moderate to severe (that is, extensive or disabling) plaque Ps (psoriasis vulgaris) with either of the following (AAD 2019):

1. Plaque Ps (psoriasis vulgaris) involving greater than three percent (3%) body surface area (BSA); **OR**
2. Plaque Ps (psoriasis vulgaris) involving less than or equal to three percent (3%) BSA involving sensitive areas or areas that significantly impact daily function (such as fingernails, palms, soles of feet, head/neck, or genitalia);

AND

B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to phototherapy or other systemic therapy (such as acitretin, cyclosporine, or methotrexate);

AND

C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to TWO (2) preferred biologic agents [Current preferred biologics include – Cosentyx (secukinumab), Enbrel (etanercept), Humira (adalimumab)] unless the following criteria are met:

1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

VIII. Non-infectious uveitis (UV) when each of the following criteria are met:

A. Individual has chronic, recurrent, treatment-refractory or vision-threatening disease; **AND**

B. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy [such as corticosteroids or immunosuppressants (azathioprine, cyclosporine, or methotrexate)];

AND

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	NA

- C. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to the preferred biologic agent [Current preferred biologic includes – Humira (adalimumab),] unless the following criteria are met:
1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

- IX. Hidradenitis suppurativa (HS) when each of the following criteria are met:
- A. Individual is 12 years of age or older; **AND**
 - B. Individual has moderate to severe hidradenitis suppurativa (Hurley stage II or Hurley stage III disease); **AND**
 - C. Individual has had an inadequate response to, is intolerant of, or has a contraindication to conventional therapy (such as oral antibiotics);
- AND**
- D. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to the preferred biologic agent [Current preferred biologic includes – Humira (adalimumab)] unless the following criteria are met:
 1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd);

OR

- X. Sarcoidosis when each of the following criteria are met (Sweiss 2014):
- A. Individual is 18 years of age or older; **AND**
 - B. Individual has chronic, progressive, treatment-refractory disease; **AND**
 - C. Individual has had an inadequate response to, is intolerant of, or has a contraindication to systemic corticosteroids; **AND**
 - D. Individual has had an inadequate response to, is intolerant of, or has a contraindication to nonbiologic disease modifying anti-rheumatic agents (DMARDs) (such as methotrexate or azathioprine);
- AND**
- E. Individual has had a trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response or intolerance to the preferred biologic agent [Current preferred biologic includes – Humira (adalimumab)] unless the following criteria are met:
 1. Individual has been receiving and is maintained on a stable dose of Hadlima (adalimumab-bwwd).

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	NA

Requests for Hadlima (adalimumab-bwwd) may **not** be approved for the following:

- I. All other indications not included above; **OR**
- II. In combination with other TNF antagonists, apremilast, JAK inhibitors or other biologic drugs (such as, abatacept, anakinra, vedolizumab); **OR**
- III. Tuberculosis, other active serious infections, or a history of recurrent infections; **OR**
- IV. Individual has not had a tuberculin skin test (TST), or a Center for Disease Control (CDC-) and Prevention -recommended equivalent, to evaluate for latent tuberculosis prior to initiating adalimumab-adbm.

Note:

TNFi have black box warnings for serious infections and malignancy. Individuals treated with TNFi are at increased risk for developing serious infections that may lead to hospitalization or death. Most individuals who developed these infections were taking concomitant immunosuppressants such as methotrexate or corticosteroids. TNFi should be discontinued if an individual develops a serious infection or sepsis. Individuals should be tested for latent tuberculosis (TB) before TNFi use and during therapy. Treatment for latent TB should be initiated prior to TNFi use. Risks and benefits of TNFi should be carefully considered prior to initiation of therapy in individuals with chronic or recurrent infection. Lymphoma and other malignancies have been reported in children and adolescents treated with TNFi. Postmarketing cases of hepatosplenic T-cell lymphoma (HSTCL) have been reported in individuals treated with TNFi. Almost all cases had received treatment with azathioprine or 6-mercaptopurine concomitantly with a TNFi at or prior to diagnosis. It is uncertain whether HSTCL is related to the use of a TNFi or a TNFi in combination with these other immunosuppressants.

Key References:

1. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2018. URL: <http://www.clinicalpharmacology.com>. Updated periodically.
2. Brunner HI, Ruperto N, Tzaribachev N, et al. Subcutaneous golimumab for children with active polyarticular-course juvenile idiopathic arthritis: results of a multicentre, double-blind, randomised-withdrawal trial. *Ann Rheum Dis*. 2018; 77(1):21-29.
3. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: September 14, 2018.
4. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
5. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2018; Updated periodically.
6. NCCN Drugs & Biologics Compendium (NCCN Compendium®) 2016 National Comprehensive Cancer Network, Inc. Available at: NCCN.org. Updated periodically. Accessed on: September 14, 2018.
7. Singh JA, Saag KG, Bridges SL et al. 2015 American College of Rheumatology Guideline for the treatment of rheumatoid arthritis. *Arthritis Rheum*. 2016;68:1-26.
8. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019; 80: 1029-72.
9. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheum*. 2019; 71(1): 5-32.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability							
Market	DC	GA	KY	MD	NJ	NY	WA
Applicable	X	X	X	X	X	X	NA

10. American Gastroenterological Association. Identification, assessment and initial medical treatment of ulcerative colitis Clinical Care Pathway. Available at <https://gastro.org/guidelines/ibd-and-bowel-disorders>. Accessed on: September 14, 2018.
11. American Gastroenterological Association. Identification, assessment and initial medical treatment of Crohn's disease Clinical Care Pathway. Available at <https://gastro.org/guidelines/ibd-and-bowel-disorders>. Accessed on: September 14, 2018.
12. Lichtenstein GR, Loftus EV, Isaacs KL et al. 2018 American College of Gastroenterology Guideline for the management of Crohn's disease in adults. *Am J Gastroenterol* 2018; 113:481–517.
13. Rubin DT, Ananthakrishnan AN, Siegel CA et al. American College of Gastroenterology Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol* 2019; 114:384-413.
14. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/ Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol.* 2019; 71(10):1599-1613.
15. Ringold S, Weiss PF, Beukelman T, et al. 2013 Update of the 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: recommendations for the medical therapy of children with systemic juvenile idiopathic arthritis and tuberculosis screening among children receiving biologic medications. *Arthritis Rheum.* 2013; 65(10):2499-2512.
16. Ringold S, Angeles-Han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. *Arthritis Rheum.* 2019; 71(6):846-863.
17. Beukelman T, Patkar NM, Saag KG, et al. 2011 American College of Rheumatology recommendations for the treatment of juvenile idiopathic arthritis: initiation and safety monitoring of therapeutic agents for the treatment of arthritis and systemic features. *Arthritis Care & Research.* 2011; 63(4):465-482.
18. Levy-Clarke G, Jabs DA, Read RW, et al. Expert panel recommendations for the use of anti-tumor necrosis factor biologic agents in patients with ocular inflammatory disorders; American Uveitis Society subcommittee. *Ophthalmology.* 2014; 121(3):785-796.
19. Baughman RP, Drent M, et al. Infliximab therapy in patients with chronic sarcoidosis and pulmonary involvement. *Am J Respir Crit Care Med.* 2006; 174:795-802.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.