

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	NA	X	X	X	X	X	NA	NA	X

\*FHK- Florida Healthy Kids

## Samsca (tolvaptan)

Override(s)	Approval Duration
Prior Authorization Quantity Limit	30 days* *Therapy duration is limited to 30 days to limit hepatic injury risk associated with medication use.

Medications	Quantity Limit
Samsca (tolvaptan) tablets	May be subject to quantity limit

### APPROVAL CRITERIA

Requests for Samsca (tolvaptan) may be approved if the following criteria are met:

- I. Individual has a diagnosis of clinically significant hypervolemic or euvolemic hyponatremia; **AND**
- II. Individual is being initiated or re-initiated on therapy in a hospital setting; **AND**
- III. Individual has a serum sodium of less than 125 mEq/L or less marked hyponatremia that is symptomatic and has resisted correction with fluid restriction.

Samsca (tolvaptan) may not be approved for the following:

- I. Individual requires an acute, urgent need to raise serum sodium; **OR**
- II. Individual is unable to sense or appropriately respond to thirst; **OR**
- III. Individual has a diagnosis of hypovolemic hyponatremia; **OR**
- IV. Individual is anuric; **OR**
- V. Individual has underlying liver disease, including cirrhosis; **OR**
- VI. Individual is currently receiving a strong CYP 3A inhibitor (such as, clarithromycin, ketoconazole, itraconazole, ritonavir, indinavir, nelfinavir, saquinavir, nefazodone, and telithromycin).

### **Notes:**

1. It has not been established that raising serum sodium with Samsca (tolvaptan) provides a symptomatic benefit.
2. Samsca (tolvaptan) has black box warnings for the need to initiate and re-initiate in a hospital and monitoring of serum sodium. Therapy should be initiated or re-initiated only in a hospital setting where serum sodium can be monitored to prevent too-rapid in

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	X	X	NA	NA	X	NA	X	X	X	X	X	NA	NA	X

\*FHK- Florida Healthy Kids

correction of hyponatremia (greater than 12 mEq/L/24 hours). Too rapid correction of hyponatremia can cause osmotic demyelination resulting in dysarthria, mutism, dysphagia, lethargy, affective changes, spastic quadriparesis, seizures, coma and death. Slower rates of sodium correction may be advisable in susceptible individuals, including those with severe malnutrition or alcoholism.

State Specific Mandates		
State name	Date effective	Mandate details (including specific bill if applicable)
N/A	N/A	N/A

**Key References:**

Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.: 2016. URL: <http://www.clinicalpharmacology.com>. Updated periodically.

DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>.

DrugPoints® System (electronic version). Truven Health Analytics, Greenwood Village, CO. Updated periodically.

Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2016; Updated periodically.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.