

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

## Zepatier (elbasvir/grazoprevir)

Override(s)	Approval Duration
Prior Authorization Quantity Limit	Based on Genotype, Treatment status, Cirrhosis status, NS5A Resistant-associated Polymorphism status, or Prior Virologic Response status

Medications	Quantity Limit
Zepatier (elbasvir/grazoprevir)	1 tablet per day

### APPROVAL DURATION

Genotype and Status (HCV mono-infected or HCV/HIV-1 co-infected <sup>a</sup> )	Associated Treatment Regimen	Total Approval Duration of Zepatier
Genotype 1b (treatment-naïve or dual P/R <sup>2b</sup> treatment-experienced, with compensated cirrhosis or without cirrhosis)	Zepatier	12 weeks
Genotype 1b (triple <sup>2d</sup> treatment-experienced, with compensated cirrhosis or without cirrhosis)	Zepatier + RBV	12 weeks
Genotype 1a (treatment-naïve or dual P/R <sup>2b</sup> treatment-experienced, with compensated cirrhosis, or without cirrhosis without baseline NS5A resistant-associated polymorphism)	Zepatier	12 weeks
Genotype 1a (treatment-naïve or dual P/R <sup>2b</sup> treatment-experienced, with compensated cirrhosis or without cirrhosis, with baseline NS5A resistant-associated polymorphism)	Zepatier + RBV	16 weeks
Genotype 1a (triple <sup>2d</sup> treatment-experienced, with compensated	Zepatier + RBV	12 weeks

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

cirrhosis or without cirrhosis, without baseline NS5A resistant-associated polymorphism)		
Genotype 1a (triple <sup>2d</sup> treatment-experienced, with compensated cirrhosis or without cirrhosis, with baseline NS5A resistant-associated polymorphism)	Zepatier + RBV	16 weeks
Genotype 3 (dual P/R <sup>2b</sup> treatment-experienced, with compensated cirrhosis)	Zepatier + Sovaldi	12 weeks
Genotype 4 (treatment-naïve, with compensated cirrhosis or without cirrhosis)	Zepatier	12 weeks
Genotype 4 (dual P/R <sup>2b</sup> treatment-experienced with virologic relapse, with compensated cirrhosis or without cirrhosis)	Zepatier	12 weeks
Genotype 4 (dual P/R <sup>2b</sup> treatment-experienced with on- treatment virologic failure <sup>†</sup> , with compensated cirrhosis or without cirrhosis)	Zepatier + RBV	16 weeks

<sup>†</sup>The September 2017 AASLD/IDSA treatment guidance defines on-treatment virologic failure as experiencing failure to suppress or breakthrough during treatment.

## **APPROVAL CRITERIA**

Requests for Zepatier (elbasvir/grazoprevir) may be approved if the following criteria are met:

- I. Individual is 18 years of age or older; **AND**
- II. Documentation is provided for a diagnosis of chronic hepatitis C (CHC) infection<sup>a</sup>, which includes genotype and a positive HCV RNA result to confirm diagnosis (AASLD/IDSA 2017, CDC 2013); **AND**
- III. Individual has received baseline evaluation for liver fibrosis to guide appropriate therapy; **AND**
- IV. Individual does not have a short life expectancy (less than 12 months owing to non-liver

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

related comorbid conditions) that cannot be remediated by treating HCV, by transplantation or other directed therapy (AASLD/IDSA 2017); **AND**

- V. Individual has compensated<sup>1</sup> liver disease (with or without cirrhosis); **AND**
- VI. If Genotype 1a subtype is present, a copy of the baseline NS5A resistant-associated polymorphism test result is provided;

**AND**

VII. Individual is using in **one** of the following antiviral treatment regimens (AASLD/IDSA 2017):

A. As monotherapy for **one** of the following:

- 1. Individual is treatment-naïve or dual P/R<sup>2b</sup> treatment-experienced, with compensated<sup>1</sup> cirrhosis or without cirrhosis, and Genotype 1b or Genotype 1a without a baseline NS5A resistant-associated polymorphism at amino acid positions M28, Q30, L31, and Y93; **AND**
- 2. Individual has had a prior trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response to Mavyret; **OR**
  - a. Individual is currently on and completing a course of therapy with Zepatier; **OR**
  - b. Documented hypersensitivity, as manifested by a severe allergic reaction to any ingredient in Mavyret which is not also in Zepatier; **OR**
  - c. Individual is concurrently using an agent that cannot be substituted with another agent or temporarily discontinued and is contraindicated or not recommended for concomitant use with the preferred regimen or regimens;

**OR**

- 3. Individual is treatment-naïve, with compensated<sup>1</sup> cirrhosis or without cirrhosis, and Genotype 4; **OR**
- 4. Individual is a dual P/R<sup>2b</sup> treatment-experienced virologic relapser, with compensated<sup>1</sup> cirrhosis or without cirrhosis, and Genotype 4;

**AND**

- 5. Individual has had a prior trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response to Epclusa **OR** Mavyret; **OR**
  - a. Individual is currently on and completing a course of therapy with Zepatier; **OR**
  - b. Documented hypersensitivity, as manifested by a severe allergic reaction to any ingredient in Epclusa **OR** Mavyret which is not also in Zepatier; **OR**

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

- c. Individual is concurrently using an agent that cannot be substituted with another agent or temporarily discontinued and is contraindicated or not recommended for concomitant use with the preferred regimen or regimens;

**OR**

B. In combination with ribavirin for **one** of the following:

1. Individual is treatment-naïve, dual P/R<sup>2b</sup>, or triple<sup>2d</sup> treatment-experienced, with compensated<sup>1</sup> cirrhosis or without cirrhosis, and Genotype 1a with a baseline NS5A resistant-associated polymorphism at amino acid positions M28, Q30, L31, and/or Y93; **OR**
2. Individual is triple<sup>2d</sup> treatment-experienced, with compensated<sup>1</sup> cirrhosis or without cirrhosis, and Genotype 1b or 1a without a baseline NS5A resistant-associated polymorphism at amino acid positions M28, Q30, L31, and Y93;

**AND**

3. Individual has had a prior trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response to Mavyret;

**OR**

- a. Individual is currently on and completing a course of therapy with Zepatier; **OR**
- b. Documented hypersensitivity, as manifested by a severe allergic reaction to any ingredient in Mavyret which is not also in Zepatier; **OR**
- c. Individual is concurrently using an agent that cannot be substituted with another agent or temporarily discontinued and is contraindicated or not recommended for concomitant use with the preferred regimen or regimens;

**OR**

4. Individual is a dual P/R<sup>2b</sup> treatment-experienced with prior on-treatment virologic failure, with compensated<sup>1</sup> cirrhosis or without cirrhosis, and Genotype 4; **AND**
5. Individual has had a prior trial (medication samples/coupons/discount cards are excluded from consideration as a trial) and inadequate response to Epclusa OR Mavyret; **OR**
  - a. Individual is currently on and completing a course of therapy with Zepatier; **OR**
  - b. Documented hypersensitivity, as manifested by a severe allergic reaction to any ingredient in Epclusa OR Mavyret which is not also in Zepatier; **OR**
  - c. Individual is concurrently using an agent that cannot be substituted with another agent or temporarily discontinued and is contraindicated or not

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

recommended for concomitant use with the preferred regimen or regimens;

**OR**

C. In combination with sofosbuvir for dual P/R<sup>2b</sup> treatment-experienced individuals, with compensated cirrhosis<sup>1</sup>, and Genotype 3.

Zepatier (elbasvir/grazoprevir) may not be approved for the following:

- I. Individual has decompensated<sup>1</sup> cirrhosis; **OR**
- II. Individual is requesting in concurrent therapy with contraindicated or not recommended agents, such as but not limited to the following: strong cytochrome (CYP) 3A4 inducers (such as but not limited to, efavirenz, phenytoin, phenobarbital, carbamazepine, St John's Wort, rifampin, rifabutin, rifapentine), organic anion transporting peptide (OATP) 1B1/1B3 inhibitors (such as but not limited to, atazanavir, darunavir, lopinavir, saquinavir, tipranavir, cyclosporine, eltrombopag, gemfibrozil, ritonavir-containing regimens), cobicistat-containing regimens, nevirapine, nafcillin, etravirine, modafinil, bosentan, oral ketoconazole; **OR**
- III. Individual is using in combination with a regimen containing another NS3/4A<sup>2c</sup> protease inhibitor; **OR**
- IV. Individual is using in combination with a regimen containing another NS5A<sup>2a</sup> inhibitor; **OR**
- V. Individual is using in combination with a regimen containing a NS5B polymerase inhibitor other than sofosbuvir (such as dasabuvir); **OR**
- VI. Individual is requesting the regimen for re-treatment and either failed to achieve a SVR (defined as a lower limit HCV RNA of 25 IU/mL) or relapsed after achieving a SVR during a prior successfully completed treatment regimen consisting of a NS5B polymerase inhibitor (such as dasabuvir or sofosbuvir) or an NS5A<sup>2a</sup> inhibitor.

**Notes:**

<sup>a</sup>Per label, Zepatier (grazoprevir/elbasvir) may be used in individuals who are co-infected with human immunodeficiency virus-1 (HIV-1).

**1. Compensated Liver Disease:**

According to the American Association for the Study of Liver Diseases (AASLD/IDSA 2017), the specific criteria for compensated liver disease include all of the following: a total bilirubin; serum albumin; prothrombin time/INR; presence of ascites; and presence of hepatic encephalopathy. However, these criteria do not establish a comprehensive definition of compensated liver disease. The AASLD guidance refers to compensated liver disease as Class A based on the Child Pugh-Turcotte (CPT) classification scoring system.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

Moderate to Severe (Decompensated) Liver Disease:

The AASLD guidance refers to decompensated (moderate to severe) liver disease as Class B or C based on the Child-Pugh Turcotte (CPT) classification scoring system.

### Child Pugh Classification (AASLD/IDSA 2017)

Parameters			
Points Assigned	1 point	2 points	3 points
Total Bilirubin (µmol/L)	<34	34-50	>50
Serum Albumin (g/L)	>35	28-35	<28
Prothrombin time/INR	<1.7	1.71-2.30	>2.30
Ascites	None	Mild	Moderate to Severe
Hepatic Encephalopathy	None	Grade I-II (or suppressed with medication)	Grade III-IV (or refractory)

### Child Pugh Score Interpretation (AASLD/IDSA 2017)

<b>Class A</b>	5-6 points	Well compensated liver disease
<b>Class B</b>	7-9 points	Significant functional compromise (moderate hepatic impairment)
<b>Class C</b>	10-15 points	Uncompensated liver disease (severe hepatic impairment)

#### 2. Past Treatment Exposure Definitions (AASLD/IDSA 2017):

- a. NS5A Inhibitor: includes daclatasvir, ledipasvir, elbasvir, ombitasvir, pibrentasvir, or velpatasvir-containing regimens
- b. P/R: includes peginterferon (or non-pegylated interferon) ± ribavirin
- c. NS3/4A Protease Inhibitor: includes simeprevir, grazoprevir, paritaprevir, glecaprevir, and voxilaprevir-containing regimens
- d. Triple therapy: includes NS3 protease inhibitor (simeprevir, boceprevir or telaprevir) plus peginterferon and ribavirin
- e. Direct Acting Antiviral (DAA): includes NS5A inhibitors, NS3/4A protease inhibitors, and NS5B polymerase inhibitors (sofosbuvir, dasabuvir)

#### 3. Chronic Kidney Disease (CKD) Definitions (AASLD/IDSA 2017):

Severe CKD (Stage 4): eGFR 15-29 mL/min

End-Stage CKD (Stage 5): eGFR < 15 mL/min

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

4. **Metavir Scoring Systems for Fibrosis Staging (AASLD 2009):**

Stage (F)	
0	No fibrosis
1	Periportal fibrotic expansion
2	Periportal septae 1 (septum)
3	Porto-central septae
4	Cirrhosis

5. Hepatitis C virus (HCV) direct acting antiviral (DAA) agents have a black box warning for risk of hepatitis B virus (HBV) reactivation in individuals with HCV-HBV co-infection. Individuals should be tested for evidence of current or prior HBV infection prior to initiation of DAA therapy. HBV reactivation has been reported in HCV/HBV co-infected individuals currently taking or previously completed DAA therapy and not concomitantly receiving HBV antiviral therapy. Some cases of HBV reactivation have led to fulminant hepatitis, hepatic failure, and death. Individuals should be monitored for hepatitis flare or HBV reactivation during and following HCV DAA therapy. Individuals should be appropriately managed for HBV infection as indicated.

State Specific Mandates		
State/Market	Date	Description
Georgia Medicaid	10/2016	Georgia has state mandated criteria; please see Georgia State Specific Criteria.
Louisiana Medicaid	2/1/2018	Louisiana has state criteria; please see Louisiana State Specific Criteria
Maryland Medicaid		Maryland has state mandated criteria; please see Maryland State Specific Criteria
Virginia Medicaid	7/1/2016	Virginia has state mandated criteria; please see Virginia State Specific Criteria.
Washington D.C.	2/1/2018	Washington D. C. has state criteria; please see Washington D. C. State Specific Criteria

**Key References:**

American Association for the Study of Liver Diseases and the Infectious Disease Society of America, in collaboration with the International Antiviral Society-USA. Recommendations for testing, managing and treating hepatitis C. Available

PAGE 7 of 8 10/29/2018

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

CRX-ALL-0295-18

Market Applicability														
Market	DC	FL & FHK	FL MMA	FL LTC	GA	KS	KY	MD	NJ	NV	NY	TN	TX	WA
Applicable	NA	X	NA	NA	NA	NA	X	NA	X	X	X	NA	NA	NA

\*FHK- Florida Healthy Kids

at <http://www.hcvguidelines.org/>. Published on: January 29, 2014. Updated on: September 21, 2017. Accessed on: January 25, 2018.

Centers for Disease Control and Prevention. Testing for HCV Infection: An Update of Guidance for Clinicians and Laboratorians. *MMWR*. 2013; 62(18):362-365. Available from: <https://www.cdc.gov/mmwr/pdf/wk/mm6218.pdf>. Accessed on: December 27, 2017.

DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed on: January 25, 2018.

European Association for the Study of the Liver. EASL Recommendations on Treatment of Hepatitis C 2016. *J Hepatol*. 2017; 66(1):153-194. Available from: [http://www.journal-of-hepatology.eu/article/S0168-8278\(16\)30489-5/pdf](http://www.journal-of-hepatology.eu/article/S0168-8278(16)30489-5/pdf). Accessed on: December 27, 2017.

U.S. Food & Drug Administration. Drugs@FDA: FDA Approved Drug Products (Package inserts). Available from: <http://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>. Accessed on: January 25, 2018.

U.S. Department of Health and Human Services AIDSinfo treatment guidelines. Concomitant use of selected antiretroviral drugs and hepatitis C virus direct-acting antiviral drugs for treatment of HCV in adults with HIV. Available at <https://aidsinfo.nih.gov/guidelines/htmltables/1/5536>. Accessed on: December 27, 2017.

DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.

Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2018; Updated periodically.

PL Detail-Document, Cytochrome P450 Drug Interactions. Pharmacist's Letter/Prescriber's Letter. May 2016.

PL Detail-Document, OATP Drug Interactions. Pharmacist's Letter/Prescriber's Letter. March 2014.

PL Detail-Document, P-glycoprotein Drug Interactions. Pharmacist's Letter/Prescriber's Letter. April 2016.

This policy does not apply to health plans or member categories that do not have pharmacy benefits, nor does it apply to Medicare. Note that market specific restrictions or transition-of-care benefit limitations may apply.

CRX-ALL-0295-18