

Preventive health guidelines

As of May 2017

What is your plan for better health?

Make this year your best year for wellness.

Your health plan will pay for recommended screenings and medically necessary tests to find disease early and will pay for routine wellness exams to help you and your family stay well. Talk with your doctor about the care that is right for you.

Your plan may not pay for all services and treatments in this guide. To learn more about what your plan pays for, see your Member Handbook or call the Member Services number on your ID card to check. You also can check www.myamerigroup.com/NJ to learn about health topics from A to Z.

The content in this guide is based in part on suggestions from these independent groups and based on state-specific requirements:

- **AAFP** – American Academy of Family Physicians
- **AAP** – American Academy of Pediatrics – Bright Futures
- **ACIP** – Advisory Committee on Immunization Practices
- **ACOG** – American Congress of Obstetricians and Gynecologists
- **ACS** – American Cancer Society
- **CDC** – Centers for Disease Control and Prevention
- **USPSTF** – U.S. Preventive Services Task Force

This guide is just for your information.

It is not meant to take the place of medical care or advice.

Use this guide to know when to set up visits with your doctor for you and your children. Ask your doctor which exams, tests and vaccines are right for you, when you should get them and how often. How you get certain diseases is not addressed in this guide.

.....

To learn more about vaccines, please see the Centers for Disease Control and Prevention (CDC) website: www.cdc.gov.



To learn more
about your plan, please see
www.myamerigroup.com/NJ.

Well-baby and child screenings

Well-baby exam – birth to 2 years*

Infants who leave the hospital less than two days (48 hours) after birth need to be seen by a doctor within two to four days after being born. You might talk to the doctor if you are a first-time parent, are having a high-risk pregnancy, or want to learn about feeding, circumcision or well-baby care. At the well-baby exam, you may get advice on your child’s safety, dental exams and care starting at age 1 year, healthy eating and development. At these exams, your baby may get vaccines and these screenings or added screenings, such as tuberculin, urine testing and/or sickle cell anemia testing, if needed.

Screenings	Age (in months)									
	Birth	1	2	4	6	9	12	15	18	24
Weight, length and head circumference (the length around the head)	At each visit, each month									
Newborn metabolic, sickle cell, and thyroid screening	Within 48 hours after birth									
Development and behavior	At each visit, each month									
Hearing	As a newborn and as your doctor suggests, at each monthly visit; referral if a deficiency									
Vision	As your doctor suggests									
Oral/dental health	Dental exams at each visit starting at age 1, then twice a year									
	Fluoride varnish on the teeth when your doctor suggests; fluoride prescription based on your drinking water									
Hemoglobin or hematocrit (blood count)	Once between 9 and 12 months, at each visit and as your doctor suggests									
Lead testing						At 12				At 24
Autism								At 18		At 24
Tuberculin	Each year and as your doctor suggests									
Urine	As your doctor suggests									

Well-child exam – ages 2½ to 10 years*

You may get advice about how to keep your child safe, how to prevent injuries, counseling to reduce the risks of getting skin cancer, good health, diet and physical activity, and development. At these well-child exams, your child may get vaccines and these screenings or added screenings such as tuberculin and urine testing, if needed.

*This guide is for people enrolled in the Amerigroup plan. Talk to your doctor if you have concerns about your health.

** Height and weight is used to find BMI. BMI is used to see if a person has the right weight for their height or is under or overweight for their height.

Age (in years)

Screenings	2½	3	4	5	6	7	8	9	10
Height, weight, body mass index (BMI)**	At each visit, each year								
Development and behavior	At each visit, each year								
Nutritional assessment	At each visit, each year								
Vision	As your doctor suggests, then at each visit, each year, starting at age 3 years								
Speech/Hearing	At each visit, each year								
Oral/dental health	Dental exams twice a year								
	Fluoride varnish on the teeth when your doctor suggests; fluoride prescription based on your drinking water								
Hemoglobin or hematocrit (blood count)	At each visit and as your doctor suggests								
Lead testing	Continues at each visit or more frequently for any elevations. Any child that has never been tested and is 72 months or less must be tested.								
Blood pressure	At age 3 and at each visit								
Dyslipidemia	As your doctor suggests								
Tuberculin	Each year and as your doctor suggests								
Urine					At 5 years				

Well-child exam – ages 11 to 18 years*

The doctor may talk to you about health and wellness issues. These include:

- Diet and physical activity
- Healthy weight
- Dental health
- Reminder to see the dentist each year
- Mental health including depression screening
- Sexual behavior and screening for sexually transmitted infections
- How to prevent injuries
- Counseling to reduce your risk of getting skin cancer; special risks you might have for cancer (such as family history) and steps you can take to reduce those risks
- Secondhand smoke
- Avoiding tobacco, alcohol and illegal drugs

At these exams, your child may get vaccines and these screenings or added screenings such as tuberculin and urine testing, if needed.

Age (in years)

Screenings	11	12	13	14	15	16	17	18
Height, weight, BMI**	At each visit, each year							
Development and behavior	At each visit, each year							
Nutritional assessment	At each visit, each year							
Blood pressure	At each visit, each year							
Vision	At each visit, each year							
Speech/Hearing	At each visit, each year							
Oral/dental health	Twice a year							
Hemoglobin or hematocrit (blood count)	As your doctor suggests; preferred at 13 years							
Chlamydia	At each visit for sexually active women age 24 and younger							
Dyslipidemia	As your doctor suggests							
Tuberculin	Each year and as your doctor suggests							
Urine	As your doctor suggests; preferred at 16 years							
Pelvic exam	Each year							

Adult screenings – women*

Well-person exam

The doctor may talk with you about health and wellness issues. These include:

- Diet and physical activity
- Family planning
- Folic acid for women who are of the age to get pregnant
- Sexual behavior and screening for sexually transmitted infections
- Screening for HIV
- Screening for Hepatitis B (HBV) if high risk
- Intimate partner violence
- How to prevent injuries
- Counseling to reduce your risk of getting skin cancer; special risks you might have for cancer (such as family history) and steps you can take to reduce those risks
- Misuse of drugs and alcohol
- Secondhand smoke
- How to stop using tobacco
- Dental health
- Mental health, including screening for depression

At this visit, you may get vaccines and these screenings:

*This guide is for people enrolled in the Amerigroup plan. Talk to your doctor if you have concerns about your health.

** Women should talk to their health care provider and choose the best age to begin screening. Women aged 50 to 74 years may have the option to screen every two years.

Screenings	Age (in years)										
	19-20	21-29	30	35	40	45	50	55	60	65 and older	
Height, weight	Each year or as your doctor suggests										
BMI	Each year or as your doctor suggests										
Blood pressure	Each year or as your doctor suggests. High measurements should be confirmed in the home setting.										
Breast cancer: Doctor exam	Every 1 to 3 years. Each year from age 39 to 65+										
Breast cancer: Mammogram	Each year from age 40 to 65+**										
Cervical cancer: ages 20-29	Every 3 years										
Cervical cancer: ages 30-65	A Pap test is recommended every 5 years. Both a Pap test and HPV test together every 5 years is preferred, but a Pap test alone can be done every 3 years.										
Cervical cancer: ages 65+	Stop screening at age 65 if last three Pap tests or last two co-tests (Pap plus HPV) within the previous 10 years were normal. If there is a history of an abnormal Pap test within the past 20 years, discuss continued screening with your doctor.										
Colorectal cancer	At age 50 and continuing until age 75 years; your doctor may suggest one or more of these test options: <ul style="list-style-type: none"> <li style="width: 50%;">■ FIT: fecal immunochemical test <li style="width: 50%;">■ Colonoscopy <li style="width: 50%;">■ FIT-DNA: multitargeted stool DNA test <li style="width: 50%;">■ CT colonography <li style="width: 50%;">■ gFOBT: guaiac-based fecal occult blood test <li style="width: 50%;">■ Flexible sigmoidoscopy 										
Chlamydia	Sexually active women ages 24 and younger										
Cholesterol	Every 5 years starting at age 40 with more screenings as your doctor suggests or for women ages 20-40 if at increased risk for coronary heart disease. Starting a medication called a statin may be recommended for some people ages 40 to 75 years who are at increased risk for cardiovascular disease.										
Glucose screening for Type 2 Diabetes	As your doctor suggests from age 40 to 70 if you are overweight or obese. Individuals with high glucose should talk to their doctor about educational counseling interventions to promote a healthful diet and physical activity.										
Hepatitis C	Screen once if born between 1945-1965										
Osteoporosis	The test to check how dense your bones are should start no later than age 65. Women at menopause should talk to their doctor about osteoporosis and have the test when at risk.										

Pregnant women*

Pregnant women should see their doctor or OB/GYN in their first three months of pregnancy for a first visit and to set up a prenatal care plan. At this visit, your doctor will check your health and the health of your baby.

Based on your past health, your doctor may want you to have these tests, screenings or vaccines:

- **Depression** – screening during or after pregnancy
- **Diabetes** – during pregnancy
- **Hematocrit/hemoglobin (blood count)**
- **Hepatitis B**
- **Hepatitis C**
- **HIV**
- **Rubella immunity** – to find out which women need the rubella vaccine after giving birth
- **Rh(D) blood type and antibody testing** – if Rh(D) negative, repeat test at 26 to 28 weeks
- **Syphilis**
- **Chlamydia**
- **Gonorrhea**
- **Group B Strep**
- **Urinalysis** – when your doctor wants it (most visits)

The doctor may talk to you about what to eat and how to be active when pregnant as well as staying away from tobacco, illegal drugs, alcohol and other substances. Your doctor will provide interventions during pregnancy and after birth to support breastfeeding, lactation supplies and counseling.



Other tests and screenings:

Some tests given alone or with other tests can be used to check the baby for health concerns. These tests are done at certain times while you are pregnant. The best test to use and the best time to do it depends on many things. These include your age as well as your medical and family history. Talk to your doctor about what these tests can tell you about your baby, the risks of the tests and which tests may be best for you.

- **Amniocentesis**
- **Chorionic villus sampling**
- **Special blood tests**
- **Ultrasound tests**, including special tests (used with blood tests during the first three months for chromosomal abnormality risk) and routine two-dimensional tests to check on the baby

Medications:

If you are high risk for a condition called preeclampsia, your doctor may recommend the use of low-dose aspirin as preventive medication.

Vaccines:


You should get the Tdap vaccine (to help protect against whooping cough), **during** your pregnancy. Other vaccines, like the flu shot, can be given before or during pregnancy, depending on whether or not it is flu season when you're pregnant.

While other vaccines may be given in special cases, it is best to get the vaccines you need before you get pregnant. Women should always check with their doctor about their own needs.

You should NOT get these vaccines while you are pregnant:

- **Measles, mumps, rubella (MMR)**
- **Varicella**

*This guide is for people enrolled in the Amerigroup plan. Talk to your doctor if you have concerns about your health.



Adult screenings – men*

Well-person exam

The doctor may talk with you about health and wellness issues. These include:

- Diet and physical activity
- Family planning
- How to prevent injuries
- Misuse of drugs and alcohol
- How to stop using tobacco
- Secondhand smoke
- Sexual behavior and screening for sexually transmitted infections
- Screening for HIV
- Screening for Hepatitis B (HBV) if high risk
- Counseling to reduce your risk of getting skin cancer; special risks you might have for cancer (such as family history) and steps you can take to reduce those risks
- Dental health
- Mental health, including screening for depression

At this visit, you may get vaccines and these screenings:

Screenings	Age in years										
	19	20	25	30	35	40	45	50	55	60	65 and older
Height, weight	Each year or as your doctor suggests										
BMI	Each year or as your doctor suggests										
Abdominal aortic aneurysm											One time for ages 65-75 for those who have ever smoked
Blood pressure	Each year or as your doctor suggests. High measurements should be confirmed in the home setting.										
Cholesterol							Every 5 years starting at age 40 with more screenings as your doctor suggests or for men ages 20-40 if at increased risk for coronary heart disease. Starting a medication called a statin may be recommended for some people ages 40 to 75 years who are at increased risk for cardiovascular disease.				
Colorectal cancer							At age 50 and continuing until age 75 years; your doctor may suggest one or more of these test options: <ul style="list-style-type: none"> ■ FIT: fecal immunochemical test ■ FIT-DNA: multitargeted stool DNA test ■ gFOBT: guaiac-based fecal occult blood test ■ Colonoscopy ■ CT colonography ■ Flexible sigmoidoscopy 				
Glucose screening for Type 2 Diabetes							As your doctor suggests from 40 to 70 if you are overweight or obese. Individuals with high glucose should talk to their doctor about educational counseling interventions to promote a healthful diet and physical activity.				
Hepatitis C							Screen once if born between 1945-1965				
Prostate cancer							If you are 50 or older, discuss with your doctor your risk of prostate cancer. Based on your history and physical exam, your doctor will recommend what is best for you.				



*This guide is for people enrolled in the Amerigroup plan. Talk to your doctor if you have concerns about your health.

Suggested vaccine schedule*

For more information about vaccinations, visit www.cdc.gov/vaccines.

*This guide is for people enrolled in the Amerigroup plan. Talk to your doctor if you have concerns about your health.

Vaccine	Age														
	Birth	1-2 months	2 months	4 months	6 months	6-18 months	12-15 months	15-18 months	19-23 months	4-6 years	11-12 years	13-18 years	19-64 years	65+ years	
Hepatitis B	✓	✓				✓									
Rotavirus (RV)			✓ 2-dose or 3-dose series												
Diphtheria, tetanus, pertussis (DTaP)			✓	✓	✓			✓		✓					
Tetanus, diphtheria, pertussis (Td/Tdap)										✓ Tdap					
Td booster													✓ Every 10 years		
Haemophilus influenza type b (Hib)			✓ 3-4 doses between 2 months to 15 months with 1st dose at 2 months, last dose at 12-15 months												
Pneumococcal conjugate (PCV)			✓	✓	✓		✓								
Inactivated polio virus (IPV)			✓	✓		✓				✓					
Influenza (flu)					✓ Suggested each year from 6 months to 65+ years of age										
Measles, mumps, rubella (MMR)							✓			✓					
Varicella (chicken pox)							✓			✓					
Hepatitis A						✓ 2-dose series between 12-23 months									
Human papillomavirus (HPV)										✓ 2-dose series					
Meningococcal										✓	✓ At age 16 MenB-FHbp: 16-23				
Pneumococcal 13-valent conjugate (PCV13)													✓ Suggested for certain individuals at risk		
Pneumococcal polysaccharide (PPSV23)													✓ Suggested for certain individuals at risk		
Zoster													✓ 1 single dose for ages 60+		

Hepatitis B – The first dose should be administered within 24 hours of birth to address births outside of the hospital. You may get an extra dose (four-dose series) at 4 months if the combination vaccine is used after the birth dose.

Rotavirus (RV) – Get two-dose or three-dose series (depends on brand of vaccine used).

Tdap (teens) – If you are 13 to 18 years of age and have not had this vaccine before, talk to your doctor about a catch-up vaccine.

Tdap (adults) – If you are 19 years of age or older and have not gotten a dose of Tdap before, you should get a single dose.

Haemophilus influenza type b (Hib) – Get three-dose or four-dose series (depends on brand of vaccine used).

Pneumococcal conjugate (PCV) – Children aged 14 months through 59 months who have received an age-appropriate series of 7-valent PCV (PCV7), get a single supplemental dose of 13-valent PCV (PCV13).

Influenza (flu) – Refer to flu.gov or cdc.gov to learn more about this vaccine. (Note: Children 6 months to 8 years of age having the vaccine for the first time should have two doses separated by four weeks.)

Measles, mumps, rubella (MMR) and varicella (chicken pox) – If you were born after 1957, you should have records of one or more doses of MMR vaccine unless you have a medical reason not to have the vaccine, or laboratory records of immunity to these diseases.

Human papillomavirus (HPV) – Eleven- to 12-year-olds receive two doses of HPV vaccine at least six months apart. Teens and young adults who start the series later, at ages 15 through 26 years, will continue to need three doses of HPV vaccine to protect against cancer-causing HPV infection.

Meningococcal – When given to healthy adolescents who are not at increased risk for meningococcal disease, 2 doses of MenB-FHbp should be administered at 0 and 6 months. If the second dose is given at an interval of less than 6 months, a third dose should be given at least 6 months after the first dose. For persons at increased risk for meningococcal disease and for use during serogroup B outbreaks, 3 doses of MenB-FHbp should be administered at 0, 1-2, and 6 months.

Pneumococcal 13-valent conjugate (PCV13)/ Pneumococcal polysaccharide (PPSV23) – Adults 65 years and older and certain adults younger than 65 who are considered at risk are recommended to receive both a PCV13 and PPSV23. Ask your doctor about the dosing recommendation that is right for you.

